

# Enyl<sup>®</sup>

Paracetamol Tablet and Suspension

## COMPOSITION

**Enyl<sup>®</sup> Tablet** : Each tablet contains Paracetamol BP 500 mg.

**Enyl<sup>®</sup> Suspension** : Each 5 ml suspension contains Paracetamol BP 120 mg.

## PHARMACOLOGY

**Enyl<sup>®</sup>** is the formulation of Paracetamol, which is one of the safest and most widely used analgesic and antipyretic. **Enyl<sup>®</sup>** produces analgesic action by elevation of the pain threshold and antipyresis through action on the hypothalamic heat regulating centre. **Enyl<sup>®</sup>** exerts significantly milder side effects and most unlikely to produce many of the serious side effects associated with aspirin and other NSAIDs.

**Enyl<sup>®</sup>** is rapidly and completely absorbed from the G.I.T. following oral administration and from rectum after rectal administration. The mean half-life of absorption from the upper small intestine is only 7 minutes. The drug is extensively metabolized in the liver and it has a plasma half-life of 1.5 to 3.0 hours. **Enyl<sup>®</sup>** is not bound to plasma proteins to any extent.

## INDICATION

Fever, common cold and influenza. Headache, toothache, earache, bodyache, myalgia, dysmenorrhoea, neuralgia and sprains. Pain of colic, back pain, post-operative pain, postpartum pain, chronic pain of cancer, inflammatory pain, and post-vaccination pain and fever of children. Rheumatism and osteoarthritic pain & stiffness of joints in fingers, hips, knees, wrists, elbows, feet, ankles and top & bottom of the spine.

## DOSAGE AND ADMINISTRATION

### Tablet :

**Adult** : 1 -2 tablets every 4 to 6 hours up to a maximum of 4 gm (8 tablets) daily.

**Children** : (6-12 years) 1/2 to 1 tablet 3 to 4 times daily.

For long term treatment, it is wise not to exceed the dose beyond 2.6 gm/day.

### Suspension

**Children** : Under 3 months: 10 mg/kg body weight (reduce to 5 mg/kg if jaundiced) 3 to 4 times daily.

3 months to below 1 year: 1/2 to 1 teaspoonful 3 to 4 times daily.

1-5 years: 1 -2 teaspoonful 3 to 4 times daily

6-12 years: 2-4 teaspoonfuls 3 to 4 times daily.

**Adults** : 4-8 teaspoonful 3 to 4 times daily.

## CONTRAINDICATION

Known sensitivity to paracetamol.

## SIDE EFFECT

Side effects are significantly mild, though haematological reactions have been reported. Pancreatitis, skin rashes, and other allergic reactions occur occasionally.

## OVER DOSE

Moderate overdose : 4-10 gm/day, Excessive overdose : More than 10 gm/day.

Symptoms of overdose:

include pallor, nausea, vomiting, anorexia & abdominal pain. Liver damage may become apparent 12-48 hours after ingestion. In severe poisoning hepatic failure may progress to encephalopathy, coma & death.

## USE IN PREGNANCY & LACTATION

Paracetamol is safe in all stages of pregnancy and lactation.

## PHARMACEUTICAL PRECAUTION

**Enyl<sup>®</sup> Tablet** and suspension : Store at a cool and dry place, protect from light and moisture

## HOW SUPPLIED

**Enyl<sup>®</sup> Tablet** : Box containing 50x10 tablets in blister pack.

**Enyl<sup>®</sup> Suspension** : 60 ml suspension in sealed cap bottle.



Manufactured by  
**EURO Pharma Ltd.**  
In Pursuit of Excellence  
Shampur, Dhaka, Bangladesh

® Regd. Trademark